Buffalo Eddy Bird Species List

Relative abundances of 43 bird species at Buffalo Eddy, Asotin County, Washington, from January 26, 1999 to December 19, 1999; 4 = abundant, 3 = common, 2 = uncommon or infrequent, 1 = rare; also shown are mean point count results and breeding status.

| Charica | Total # ^a | Birds/Count Period ^a | Density (Birds/ha) ^a | 90% Cl ^a | Total #b | % of | Abundance | Breeding |
|---------------------------------------|----------------------|------------------------------------|------------------------------------|------------------------|----------|---------------------------|---------------|---------------------|
| Species Canada goose | rotal# | Period | (Birds/na) | CI | 13 | Total ^b 7.3 | Category 4 | Status ^c |
| | | | | | | 7.3 0.6 | 2 | ı M |
| Common merganser | | | | | 1 5 | | 3 | В |
| Northern harrier | | | | | 5 | 2.8 | 3 2 | Б Т |
| Cooper's hawk ^d | | | | | 2 | 4 7 | 3 | T |
| Red-tailed hawk | | | | | 3 | 1.7 | | T |
| Golden eagle | | | | | 1 | 0.6 | 2 | В |
| American kestrel | | | | | 10 | 5.6 | 4 | |
| Chukar | | | | | 8 2 | 4.5 | 3 2 | L B |
| Gray partridge | | 0.44 | | | | 1.1 | | |
| California quail | 1 | 0.11 | | | 1 | 0.6 | 2 2 | L |
| Killdeer ^d | | | | | 0 | 5 4 | | M |
| Rock pigeon | | | | | 9 | 5.1 | 4 | L |
| Mourning dove | | | | | 2 | 1.1 | 2 | L |
| Western screech-owl ^d | | | | | | | 2 | M |
| Common nighthawkl ^d | | | | | | | 2 | T |
| Unidentified hummingbird ^d | | 0.44 | | | 4 | 0.0 | 1 | M |
| Belted kingfisher | 1 | 0.11 | | | 1 | 0.6 | 2 | T |
| Northern flicker | | | | | 2 | 1.1 | 2 | T T |
| Willow flycatcher ^d | 0 | 0.00 | 0.050 | 0.070 | 0 | 0.4 | 1 | |
| Western kingbird | 2 | 0.22 | 0.053 | 0.079 | 6 | 3.4 | 3 | B T |
| Black-billed magpie | 1 | 0.11 | | | 2 | 1.1 | 2 | |
| American crow | | | | | 18 | 10.2 | 4 | L |
| Violet-green swallow | | | | | 9 | 5.1 | 4 | L |
| Black-capped chickadee | | 0.44 | | | 1 | 0.6 | 2 | M |
| Rock wren | 1 | 0.11 | | | 5 | 2.8 | 3 | В |
| Canyon wren | | | | | 19 | 10.7 | 4 | В |
| Bewick's wren ^d | | | | | | | 1 | T |
| Winter wren ^d | | | | | | | 1 | M |
| Golden-crowned kinglet ^d | | | | | | | 1 | M |
| Ruby-crowned kinglet ^d | • | 0.07 | 0.044 | 0.074 | • | 0.4 | 1 | M |
| Yellow warbler | 6 | 0.67 | 0.314 | 0.274 | 6 | 3.4 | 3 | L |
| Yellow-rumped warbler ^d | | | | | | | 1 | M |
| MacGillivray's warbler | | | | | 1 | 0.6 | 2 | M |
| Yellow-breasted chat | _ | | | | 2 | 1.1 | 2 | L |
| Spotted towhee | 2 | 0.22 | | | 9 | 5.1 | 4 | L |
| Song sparrow | 8 | 0.89 | 0.605 | 0.515 | 16 | 9.0 | 4 | В |
| Dark-eyed junco ^d | _ | | | | _ | | 1 | M |
| Black-headed grosbeak | 1 | 0.11 | | | 2 | 1.1 | 2 | L |
| Lazuli bunting | 5 | 0.56 | 0.307 | 0.313 | 9 | 5.1 | 4 | В |
| Red-winged blackbird | | | | | 2 | 1.1 | 2 | M |
| Western meadowlark ^d | | | | | | | 1 | T |
| Brown-headed cowbird | 1 | 0.11 | | | 5 | 2.8 | 3 | T |
| American goldfinch | 1 | 0.11 | | | 7 | 4.0 | 3 | L |

 ^a Breeding season point count results (data truncated at 100 m). Sample sizes of one were too small for density analysis and were eliminated from the table.
 ^b Total number recorded and percent of total based on unlimited distance point counts.
 ^c B = Breeder; L = Likely Breeder; T = Transient; M = Migrant.
 ^d Encountered outside of point counts.